




SIDIS topics at the EIC

Charlotte Van Hulse
UAH

Kinematic coverage for DIS




Helicity structure of the nucleon: gluons





Inclusive measurements
→ access to gluon spin

High e^- purity needed, e.g. ECCE via EMCAL:
<2% π contamination for $p > 2$ GeV





Helicity structure of the nucleon: gluons





Inclusive measurements
→ access to gluon spin

High e^- purity needed, e.g. ECCE via EMCAL:
<2% π contamination for $p > 2$ GeV







Helicity structure of the nucleon: gluons





Inclusive measurements
→ access to gluon spin



High e^- purity needed, e.g. ECCE via EMCAL:
<2% π contamination for $p > 2$ GeV



Helicity structure of the nucleon via collinear SIDIS





Helicity structure of the nucleon via collinear SIDIS



$$A_{\parallel}^h(x_B, Q^2, z) = \frac{1}{P_e P_p} \frac{\frac{\overrightarrow{N}^h}{\overrightarrow{L}} - \frac{\overleftarrow{N}^h}{\overleftarrow{L}}}{\frac{\overrightarrow{N}^h}{\overrightarrow{L}} + \frac{\overleftarrow{N}^h}{\overleftarrow{L}}} (x_B, Q^2, z)$$


$$= D(y) A_1^h(x_B, Q^2, z) \quad \left(z \stackrel{\text{lab}}{=} \frac{E_h}{E_{\gamma^*}} \right)$$

Helicity structure of the nucleon via collinear SIDIS




Semi-inclusive measurements, via good hadron PID
→ access to sea-quark spin

$$\begin{aligned}
 A_{\parallel}^h(x_B, Q^2, z) &= \frac{1}{P_e P_p} \frac{\frac{\overrightarrow{N}^h}{\overrightarrow{L}} - \frac{\overleftarrow{N}^h}{\overleftarrow{L}}}{\frac{\overrightarrow{N}^h}{\overrightarrow{L}} + \frac{\overleftarrow{N}^h}{\overleftarrow{L}}} (x_B, Q^2, z) \\
 &= D(y) A_1^h(x_B, Q^2, z) \quad \left(z \stackrel{\text{lab}}{=} \frac{E_h}{E_{\gamma^*}} \right) \\
 &\propto \sum_q e_q^2 \left[\Delta q \otimes w_1 D_1^{q \rightarrow h} \right]
 \end{aligned}$$



Helicity structure of the nucleon via collinear SIDIS





Semi-inclusive measurements, via good hadron PID
→ access to sea-quark spin

$$\begin{aligned}
 A_{\parallel}^h(x_B, Q^2, z) &= \frac{1}{P_e P_p} \frac{\frac{\overrightarrow{N}^h}{\overrightarrow{L}} - \frac{\overleftarrow{N}^h}{\overleftarrow{L}}}{\frac{\overrightarrow{N}^h}{\overrightarrow{L}} + \frac{\overleftarrow{N}^h}{\overleftarrow{L}}} (x_B, Q^2, z) \\
 &= D(y) A_1^h(x_B, Q^2, z) \quad \left(z \stackrel{\text{lab}}{=} \frac{E_h}{E_{\gamma^*}} \right) \\
 &\propto \sum_q e_q^2 \left[\Delta q \otimes w_1 D_1^{q \rightarrow h} \right]
 \end{aligned}$$

Beam polarisations assumed to be 70%.

Generated and reconstructed A_1

$$D(y) = 1$$




General good agreement between reconstructed and generated asymmetry: moderate smearing.


Kinematic coverage and uncertainties

ECCE

u, d sea quarks





strange quarks



via tracking of
scattered
lepton only



Kinematic coverage and uncertainties

ECCE



via tracking of
scattered
lepton only



Helicity structure of the nucleon via SIDIS



Reweightings of Pythia MC at NLO,
using DSSV14 helicity distributions
and DSS14 pion and kaon FFs


systematic uncertainty = |generated - reconstructed|

Helicity structure of the nucleon via SIDIS




Reweightings of Pythia MC at NLO,
using DSSV14 helicity distributions
and DSS14 pion and kaon FFs

systematic uncertainty = |generated - reconstructed|




Transverse-momentum dependent SIDIS cross section for single-hadron production

$$\begin{aligned}
 \sigma^h(\phi, \phi_S) = & \sigma_{UU}^h \left\{ 1 + 2\langle \cos(\phi) \rangle_{UU}^h \cos(\phi) + 2\langle \cos(2\phi) \rangle_{UU}^h \cos(2\phi) \right. \\
 & + \lambda_l 2\langle \sin(\phi) \rangle_{LU}^h \sin(\phi) \\
 & + S_L \left[2\langle \sin(\phi) \rangle_{UL}^h \sin(\phi) + 2\langle \sin(2\phi) \rangle_{UL}^h \sin(2\phi) \right. \\
 & + \lambda_l \left(2\langle \cos(0\phi) \rangle_{LL}^h \cos(0\phi) + 2\langle \cos(\phi) \rangle_{LL}^h \cos(\phi) \right) \Big] \\
 & + S_T \left[2\langle \sin(\phi - \phi_S) \rangle_{UT}^h \sin(\phi - \phi_S) + 2\langle \sin(\phi + \phi_S) \rangle_{UT}^h \sin(\phi + \phi_S) \right. \\
 & + 2\langle \sin(3\phi - \phi_S) \rangle_{UT}^h \sin(3\phi - \phi_S) + 2\langle \sin(\phi_S) \rangle_{UT}^h \sin(\phi_S) \\
 & + 2\langle \sin(2\phi - \phi_S) \rangle_{UT}^h \sin(2\phi - \phi_S) \\
 & + \lambda_l \left(2\langle \cos(\phi - \phi_S) \rangle_{LT}^h \cos(\phi - \phi_S) \right. \\
 & + \left. \left. 2\langle \cos(\phi_S) \rangle_{LT}^h \cos(\phi_S) + 2\langle \cos(2\phi - \phi_S) \rangle_{LT}^h \cos(2\phi - \phi_S) \right) \right] \Big\}
 \end{aligned}$$



Transverse-momentum dependent SIDIS cross section for single-hadron production


$$\begin{aligned}
 \sigma^h(\phi, \phi_S) = & \sigma_{UU}^h \left\{ 1 + 2\langle \cos(\phi) \rangle_{UU}^h \cos(\phi) + 2\langle \cos(2\phi) \rangle_{UU}^h \cos(2\phi) \right. \\
 & + \lambda_l 2\langle \sin(\phi) \rangle_{LU}^h \sin(\phi) \\
 & \xleftarrow{\text{longitudinal target polarization}} + S_L \left[2\langle \sin(\phi) \rangle_{UL}^h \sin(\phi) + 2\langle \sin(2\phi) \rangle_{UL}^h \sin(2\phi) \right. \\
 & \left. + \lambda_l \left(2\langle \cos(0\phi) \rangle_{LL}^h \cos(0\phi) + 2\langle \cos(\phi) \rangle_{LL}^h \cos(\phi) \right) \right] \\
 & \xleftarrow{\text{transverse target polarization}} + S_T \left[2\langle \sin(\phi - \phi_S) \rangle_{UT}^h \sin(\phi - \phi_S) + 2\langle \sin(\phi + \phi_S) \rangle_{UT}^h \sin(\phi + \phi_S) \right. \\
 & + 2\langle \sin(3\phi - \phi_S) \rangle_{UT}^h \sin(3\phi - \phi_S) + 2\langle \sin(\phi_S) \rangle_{UT}^h \sin(\phi_S) \\
 & + 2\langle \sin(2\phi - \phi_S) \rangle_{UT}^h \sin(2\phi - \phi_S) \\
 & \xleftarrow{\text{beam polarization}} + \lambda_l \left(2\langle \cos(\phi - \phi_S) \rangle_{LT}^h \cos(\phi - \phi_S) \right. \\
 & \left. + 2\langle \cos(\phi_S) \rangle_{LT}^h \cos(\phi_S) + 2\langle \cos(2\phi - \phi_S) \rangle_{LT}^h \cos(2\phi - \phi_S) \right) \left. \right\}
 \end{aligned}$$



Transverse-momentum dependent SIDIS cross section for single-hadron production

$$\begin{aligned}
 \sigma^h(\phi, \phi_S) = & \sigma_{UU}^h \left\{ 1 + 2\langle \cos(\phi) \rangle_{UU}^h \cos(\phi) + 2\langle \cos(2\phi) \rangle_{UU}^h \cos(2\phi) \right. \\
 & + \lambda_l 2\langle \sin(\phi) \rangle_{LU}^h \sin(\phi) \\
 & \xleftarrow{\text{longitudinal target polarization}} + S_L \left[2\langle \sin(\phi) \rangle_{UL}^h \sin(\phi) + 2\langle \sin(2\phi) \rangle_{UL}^h \sin(2\phi) \right. \\
 & \left. + \lambda_l \left(2\langle \cos(0\phi) \rangle_{LL}^h \cos(0\phi) + 2\langle \cos(\phi) \rangle_{LL}^h \cos(\phi) \right) \right] \\
 & \xleftarrow{\text{transverse target polarization}} + S_T \left[2\langle \sin(\phi - \phi_S) \rangle_{UT}^h \sin(\phi - \phi_S) + 2\langle \sin(\phi + \phi_S) \rangle_{UT}^h \sin(\phi + \phi_S) \right. \\
 & + 2\langle \sin(3\phi - \phi_S) \rangle_{UT}^h \sin(3\phi - \phi_S) + 2\langle \sin(\phi_S) \rangle_{UT}^h \sin(\phi_S) \\
 & + 2\langle \sin(2\phi - \phi_S) \rangle_{UT}^h \sin(2\phi - \phi_S) \\
 & \xleftarrow{\text{beam polarization}} + \lambda_l \left(2\langle \cos(\phi - \phi_S) \rangle_{LT}^h \cos(\phi - \phi_S) \right. \\
 & \left. + 2\langle \cos(\phi_S) \rangle_{LT}^h \cos(\phi_S) + 2\langle \cos(2\phi - \phi_S) \rangle_{LT}^h \cos(2\phi - \phi_S) \right) \left. \right\}
 \end{aligned}$$

beam polarization
target polarization





Transverse-momentum dependent SIDIS cross section for single-hadron production

$$\begin{aligned}
 \sigma^h(\phi, \phi_S) = & \sigma_{UU}^h \left\{ 1 + 2\langle \cos(\phi) \rangle_{UU}^h \cos(\phi) + 2\langle \cos(2\phi) \rangle_{UU}^h \cos(2\phi) \right. \\
 & + \lambda_l 2\langle \sin(\phi) \rangle_{LU}^h \sin(\phi) \\
 & \xleftarrow{\text{longitudinal target polarization}} + S_L \left[2\langle \sin(\phi) \rangle_{UL}^h \sin(\phi) + 2\langle \sin(2\phi) \rangle_{UL}^h \sin(2\phi) \right. \\
 & \xleftarrow{\text{transverse target polarization}} + \lambda_l \left(2\langle \cos(0\phi) \rangle_{LL}^h \cos(0\phi) + 2\langle \cos(\phi) \rangle_{LL}^h \cos(\phi) \right) \\
 & + S_T \left[2\langle \sin(\phi - \phi_S) \rangle_{UT}^h \sin(\phi - \phi_S) + 2\langle \sin(\phi + \phi_S) \rangle_{UT}^h \sin(\phi + \phi_S) \right. \\
 & + 2\langle \sin(3\phi - \phi_S) \rangle_{UT}^h \sin(3\phi - \phi_S) + 2\langle \sin(\phi_S) \rangle_{UT}^h \sin(\phi_S) \\
 & + 2\langle \sin(2\phi - \phi_S) \rangle_{UT}^h \sin(2\phi - \phi_S) \\
 & \xleftarrow{\text{beam polarization}} + \lambda_l \left(2\langle \cos(\phi - \phi_S) \rangle_{LT}^h \cos(\phi - \phi_S) \right. \\
 & \left. + 2\langle \cos(\phi_S) \rangle_{LT}^h \cos(\phi_S) + 2\langle \cos(2\phi - \phi_S) \rangle_{LT}^h \cos(2\phi - \phi_S) \right) \left. \right\}
 \end{aligned}$$

beam polarization target polarization

leading twist





Transverse-momentum dependent SIDIS cross section for single-hadron production

$$\begin{aligned}
 \sigma^h(\phi, \phi_S) = & \sigma_{UU}^h \left\{ 1 + 2\langle \cos(\phi) \rangle_{UU}^h \cos(\phi) + 2\langle \cos(2\phi) \rangle_{UU}^h \cos(2\phi) \right. \\
 & + \lambda_l 2\langle \sin(\phi) \rangle_{LU}^h \sin(\phi) \\
 & \xleftarrow{\text{longitudinal target polarization}} + S_L \left[2\langle \sin(\phi) \rangle_{UL}^h \sin(\phi) + 2\langle \sin(2\phi) \rangle_{UL}^h \sin(2\phi) \right. \\
 & \xleftarrow{\text{transverse target polarization}} + \lambda_l \left(2\langle \cos(0\phi) \rangle_{LL}^h \cos(0\phi) + 2\langle \cos(\phi) \rangle_{LL}^h \cos(\phi) \right) \\
 & + S_T \left[2\langle \sin(\phi - \phi_S) \rangle_{UT}^h \sin(\phi - \phi_S) + 2\langle \sin(\phi + \phi_S) \rangle_{UT}^h \sin(\phi + \phi_S) \right. \\
 & + 2\langle \sin(3\phi - \phi_S) \rangle_{UT}^h \sin(3\phi - \phi_S) + 2\langle \sin(\phi_S) \rangle_{UT}^h \sin(\phi_S) \\
 & + 2\langle \sin(2\phi - \phi_S) \rangle_{UT}^h \sin(2\phi - \phi_S) \\
 & \xleftarrow{\text{beam polarization}} + \lambda_l \left(2\langle \cos(\phi - \phi_S) \rangle_{LT}^h \cos(\phi - \phi_S) \right. \\
 & + 2\langle \cos(\phi_S) \rangle_{LT}^h \cos(\phi_S) + 2\langle \cos(2\phi - \phi_S) \rangle_{LT}^h \cos(2\phi - \phi_S) \left. \left. \right] \right\}
 \end{aligned}$$

beam polarization target polarization

sub-leading twist




Transverse-momentum dependent SIDIS cross section for single-hadron production

$$\begin{aligned}
 \sigma^h(\phi, \phi_S) = & \sigma_{UU}^h \left\{ 1 + 2\langle \cos(\phi) \rangle_{UU}^h \cos(\phi) + 2\langle \cos(2\phi) \rangle_{UU}^h \cos(2\phi) \right. \\
 & + \lambda_l 2\langle \sin(\phi) \rangle_{LU}^h \sin(\phi) \\
 & \xleftarrow{\text{longitudinal target polarization}} + S_L \left[2\langle \sin(\phi) \rangle_{UL}^h \sin(\phi) + 2\langle \sin(2\phi) \rangle_{UL}^h \sin(2\phi) \right. \\
 & + \lambda_l \left(2\langle \cos(0\phi) \rangle_{LL}^h \cos(0\phi) + 2\langle \cos(\phi) \rangle_{LL}^h \cos(\phi) \right) \\
 & \xleftarrow{\text{transverse target polarization}} + S_T \left[2\langle \sin(\phi - \phi_S) \rangle_{UT}^h \sin(\phi - \phi_S) + 2\langle \sin(\phi + \phi_S) \rangle_{UT}^h \sin(\phi + \phi_S) \right. \\
 & + 2\langle \sin(3\phi - \phi_S) \rangle_{UT}^h \sin(3\phi - \phi_S) + 2\langle \sin(\phi_S) \rangle_{UT}^h \sin(\phi_S) \\
 & + 2\langle \sin(2\phi - \phi_S) \rangle_{UT}^h \sin(2\phi - \phi_S) \\
 & \xleftarrow{\text{beam polarization}} + \lambda_l \left(2\langle \cos(\phi - \phi_S) \rangle_{LT}^h \cos(\phi - \phi_S) \right. \\
 & + \left. \left. 2\langle \cos(\phi_S) \rangle_{LT}^h \cos(\phi_S) + 2\langle \cos(2\phi - \phi_S) \rangle_{LT}^h \cos(2\phi - \phi_S) \right) \right] \Big\}
 \end{aligned}$$

beam polarization
target polarization

presented here



TMD PDFs and fragmentation functions (FFs)

Azimuthal amplitudes related to structure functions F_{XY} :


$$2\langle \sin(\phi + \phi_S) \rangle_{UT}^h = \epsilon F_{UT}^{\sin(\phi + \phi_S)}$$

TMD PDFs and fragmentation functions (FFs)

Azimuthal amplitudes related to structure functions F_{XY} :

$$2\langle \sin(\phi + \phi_S) \rangle_{UT}^h = \epsilon F_{UT}^{\sin(\phi + \phi_S)}$$

$$F_{XY} \propto \mathcal{C} [\text{TMD PDF}(x, k_{\perp}) \times \text{TMD FF}(z, p_{\perp})]$$





TMD PDFs and fragmentation functions (FFs)

Azimuthal amplitudes related to structure functions F_{XY} :

$$2\langle \sin(\phi + \phi_S) \rangle_{UT}^h = \epsilon F_{UT}^{\sin(\phi + \phi_S)}$$

$$F_{XY} \propto \mathcal{C} [\text{TMD PDF}(x, k_{\perp}) \times \text{TMD FF}(z, p_{\perp})]$$




TMD PDFs and fragmentation functions (FFs)

Azimuthal amplitudes related to structure functions F_{XY} :

$$2\langle \sin(\phi + \phi_S) \rangle_{UT}^h = \epsilon F_{UT}^{\sin(\phi + \phi_S)}$$

$$F_{XY} \propto \mathcal{C} [\text{TMD PDF}(x, k_\perp) \times \text{TMD FF}(z, p_\perp)]$$

		quark polarization		
		U	L	T
nucleon polarization	U	f_1		h_1^\perp
	L		g_{1L}	h_{1L}^\perp
	T	f_{1T}^\perp	g_{1T}^\perp	$h_{1T} h_{1T}^\perp$




TMD PDFs and fragmentation functions (FFs)

Azimuthal amplitudes related to structure functions F_{XY} :


$$2\langle \sin(\phi + \phi_S) \rangle_{UT}^h = \epsilon F_{UT}^{\sin(\phi + \phi_S)}$$

$$F_{XY} \propto \mathcal{C} [\text{TMD PDF}(x, k_\perp) \times \text{TMD FF}(z, p_\perp)]$$



		quark polarization		
		U	L	T
nucleon polarization	U	f_1		h_1^\perp
	L		g_{1L}	h_{1L}^\perp
	T	f_{1T}^\perp	g_{1T}^\perp	$h_{1T} h_{1T}^\perp$

		quark polarization		
		U	L	T
hadron polarization	U	D_1		H_1^\perp




TMD PDFs and fragmentation functions (FFs)

Azimuthal amplitudes related to structure functions F_{XY} :


$$2\langle \sin(\phi + \phi_S) \rangle_{UT}^h = \epsilon F_{UT}^{\sin(\phi + \phi_S)}$$

$$F_{XY} \propto \mathcal{C} [\text{TMD PDF}(x, k_\perp) \times \text{TMD FF}(z, p_\perp)]$$



		quark polarization		
		U	L	T
nucleon polarization	U	f_1		h_1^\perp
	L		g_{1L}	h_{1L}^\perp
	T	f_{1T}^\perp	g_{1T}^\perp	$h_{1T} h_{1T}^\perp$

		quark polarization		
		U	L	T
hadron polarization	U	D_1		H_1^\perp
	L			




TMD PDFs and fragmentation functions (FFs)

Azimuthal amplitudes related to structure functions F_{XY} :


$$2\langle \sin(\phi + \phi_S) \rangle_{UT}^h = \epsilon F_{UT}^{\sin(\phi + \phi_S)}$$

$$F_{XY} \propto \mathcal{C} [\text{TMD PDF}(x, k_\perp) \times \text{TMD FF}(z, p_\perp)]$$




		quark polarization		
		U	L	T
nucleon polarization	U	f_1		h_1^\perp
	L		g_{1L}	h_{1L}^\perp
	T	f_{1T}^\perp	g_{1T}^\perp	$h_{1T} h_{1T}^\perp$

		quark polarization		
		U	L	T
hadron polarization	U	D_1		H_1^\perp



Spin-independent TMD PDF




Fit:
A. Bacchetta et al.,
JHEP 06 (2017) 081,
JHEP 06 (2019) 051 (erratum)

EIC uncertainties dominated
by assumed
3% point-to-point uncorrelated uncertainty
3% scale uncertainty

Theory uncertainties dominated by
TMD evolution.

Spin-independent TMD PDF

Large lever-arm in Q^2 over large x range
 → Q^2 evolution of TMD PDF





Fit:
 A. Bacchetta et al.,
 JHEP 06 (2017) 081,
 JHEP 06 (2019) 051 (erratum)

EIC uncertainties dominated
 by assumed
 3% point-to-point uncorrelated uncertainty
 3% scale uncertainty

Theory uncertainties dominated by
 TMD evolution.

Spin-independent TMD PDF: impact of EIC



DIS variables via scattered lepton

$$\begin{aligned} Q^2 &> 1 \text{ GeV}^2 \\ 0.01 < y < 0.95 \\ W^2 &> 10 \text{ GeV}^2 \end{aligned}$$

$$\begin{aligned} 5 \times 41 \text{ GeV}^2 \\ 10 \times 100 \text{ GeV}^2 \\ 18 \times 100 \text{ GeV}^2 \\ 18 \times 275 \text{ GeV}^2 \end{aligned}$$

$$\mathcal{L} = 10 \text{ fb}^{-1} \text{ for each collision energy}$$

$$\text{systematic uncertainty} = |\text{generated} - \text{reconstructed}|$$


Fit: I. Scimemi, A. Vladimirov
JHEP, 06:137, 2020

TMD evolution of the Sivers TMD PDF



$$\mathcal{C}[f_{1T}^\perp \times D_1^{q \rightarrow h}]$$

M. Bury et al., JHEP, 05:151, 2021.




Decrease of asymmetry with increasing $Q^2 \rightarrow$ need high precision ($<1\%$) to measure asymmetry at high Q^2

Generated and reconstructed Sivers asymmetry

Reweighting of Pythia at LO using Sivers extraction from existing SIDIS and e^+e^- data

M. Anselmino et al., Eur. Phys. J. A 39:89-100, 2009



General good agreement between reconstructed and generated asymmetry: moderate smearing.

Uncertainties Sivers asymmetry




$$\mathcal{C}[f_{1T}^\perp \times D_1^{q \rightarrow h}]$$

Beam polarisations assumed to be 70%.


systematic uncertainty=
|generated - reconstructed|

Additionally: 3% scale uncertainty




- Low x and Q^2 : small statistical uncertainty. High precision is needed since asymmetry at low x and Q^2 well below 1%.
- For not too large z and P_T , statistical uncertainty well below 1%.
- Systematic uncertainties increase with z and P_T : likely because of higher smearing effects.

Uncertainties Sivers asymmetry



Q^2 dependence of the Sivers asymmetry





Intermediate and high x : good coverage in Q^2 ,
with complementarity in coverage at different COM energies.

Sivers TMD PDF: impact of EIC

Parametrisation from
M. Bury et al., JHEP, 05:151, 2021

$Q=2\text{ GeV}$



ECCE



DIS variables via scattered lepton

$$\begin{array}{ll}
 Q^2 > 1 \text{ GeV}^2 & 5 \times 41 \text{ GeV}^2 \\
 0.01 < y < 0.95 & 10 \times 100 \text{ GeV}^2 \\
 & 18 \times 100 \text{ GeV}^2 \\
 W^2 > 10 \text{ GeV}^2 & 18 \times 275 \text{ GeV}^2
 \end{array}$$

$\mathcal{L} = 10 \text{ fb}^{-1}$ for each collision energy

Transversity: impact of EIC

Parametrisation from J. Cammarota, Phys. Rev. D 102(5):054002, 2020.



DIS variables via scattered lepton

$$Q^2 > 1 \text{ GeV}^2$$

$$0.01 < y < 0.95$$

$$W^2 > 10 \text{ GeV}^2$$

$$5 \times 41 \text{ GeV}^2$$

$$10 \times 100 \text{ GeV}^2$$


$$18 \times 100 \text{ GeV}^2$$

$$18 \times 275 \text{ GeV}^2$$

$$\mathcal{L} = 10 \text{ fb}^{-1} \text{ for each collision energy}$$

Back-to-back dihadron production in eA

- Access to the Weizsäcker-Williams gluon distribution
- Sensitive to saturation effects
- Complementarity region covered by dihadron and jet production



Summary and outlook

Semi-inclusive measurements at EIC provide access to a range of information:

- Helicity distributions of sea and valence quarks
- 3D (spin-dependent) momentum structure of the nucleon, and study of TMD evolution
- Probe gluon saturation

Full GEANT simulations have been performed by ATHENA and ECCE, which demonstrate the EIC capabilities to perform measurements with a large kinematic coverage and with high precision.


This is needed in order to allow for a precise determination of the various distributions.

Next steps:


- reference design optimisation and consolidation phase, with joints efforts from ATHENA and ECCE.
- towards the formation of a new detector 1 collaboration.

Back up

Sivers: statistical and systematic uncertainties



Sivers: statistical and systematic uncertainties



Sivers: Q^2 coverage

